

On UV/IR Mixing via Seiberg-Witten Map for Noncommutative QED

Matti Raasakka^{1,†} and Anca Tureanu^{1,2}

¹*Department of Physics, University of Helsinki*

²*Helsinki Institute of Physics,*

P.O. Box 64, FIN-00014 Helsinki, Finland

`matti.raasakka@aei.mpg.de, anca.tureanu@helsinki.fi`

Abstract

We consider quantum electrodynamics in noncommutative spacetime by deriving a θ -exact Seiberg-Witten map with fermions in the fundamental representation of the gauge group as an expansion in the coupling constant. Accordingly, we demonstrate the persistence of UV/IR mixing in noncommutative QED with charged fermions via Seiberg-Witten map, extending the results of Schupp and You [1].

1 Introduction

The construction of renormalizable quantum field theories in noncommutative spacetime endowed with canonical coordinate commutation relations $[\hat{x}^\mu, \hat{x}^\nu] = i\theta^{\mu\nu}$ is a long-standing problem, the solution of which is necessary for the calculation of testable predictions for these theories. They are expected to give hints of the underlying quantum structure of spacetime, in particular, due to their appearance in string theory [2] and in semi-classical situations, where principles of quantum field theory and general relativity are combined [3]. Arguably, the most serious obstacle for the formulation of these noncommutative quantum field theories is the so-called UV/IR mixing, giving rise to nonrenormalizable divergencies, which seem to be a generic property of any quantum field theory in noncommutative spacetime due to the inherent infinite range of nonlocality induced by the noncommutativity. Various solutions have been proposed to cure the problem. (See e.g. [4] for a review.)

[†]Present address: Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 4, D-14476 Golm, Germany.

In their seminal paper [2] on the connection between noncommutative geometry and string theory, Seiberg and Witten introduced a mapping, which relates gauge field theories in noncommutative spacetime to ordinary commutative ones, known as the Seiberg-Witten map. This mapping has virtues, since many aspects of gauge theories, such as observables and gauge fixing, are more easily understood and dealt with in the language of ordinary theories. On the other hand, it also has certain uniqueness ambiguities explored in [5, 6]. Moreover, it does not seem to affect at all some problems stemming from the noncommutativity, an example of which is the no-go theorem [7, 8], according to which fields can transform nontrivially under only two different gauge groups $U_\star(N)$.

Therefore it is interesting to study whether the Seiberg-Witten map affects the problem of UV/IR mixing. Indeed, it has been argued, for example in [9]¹, that the mixing of UV and IR sectors of noncommutative theories is absent in the Seiberg-Witten formalism. However, we suspect that this may be due to the expansion in the noncommutativity parameter matrix θ in the θ -expanded Seiberg-Witten map. In the θ -exact Seiberg-Witten map for noncommutative QED, the UV/IR mixing problem does appear, as we shall demonstrate. The same argument has been expressed by Schupp and You in [1], where they considered a noncommutative model with a gauge field coupled with a spinor field in the adjoint representation of the gauge group, and showed the existence of an IR-divergent term for the photon self-energy corrections. The adjoint representation, however, corresponds to a chargeless particle but with an electric dipole moment proportional to θ [10, 11, 12, 13], and thus in their model the interaction vanishes at the commutative limit $\theta \rightarrow 0$. Therefore the model does not correspond to a noncommutative theory of electrically charged fermions, which should reduce to the ordinary QED in the commutative limit.

In this paper, our primary goal is to extend the analysis of Ref. [1] to the case of noncommutative QED with charged fermions. We first derive a θ -exact Seiberg-Witten map for a gauge theory with a spinor field in the fundamental representation of the gauge field, corresponding to charged fermions, as an expansion in the coupling constant, and then demonstrate the persistence of UV/IR mixing in the photon self-energy corrections.

2 θ -exact Seiberg-Witten map with charged fermions

The Seiberg-Witten map, as introduced in Ref. [2], is a technique to induce a gauge orbit preserving mapping $(A_\mu, \Lambda) \mapsto (\hat{A}_\mu, \hat{\Lambda})$ between gauge fields and gauge transformation parameters in commutative and noncommutative spacetimes, respectively. The

¹In particular, in [9] it was shown that via Seiberg-Witten map the photon self-energy diagram can be renormalized up to any *finite* order in θ by shifting the nonrenormalizable terms up to the next order.

mapping has previously been realized either as an expansion in the noncommutativity parameters $\theta^{\mu\nu}$ or in the gauge field A_μ as established for an Abelian gauge field theory in [14, 15, 1] (in the respective order). However, there is a third way, namely, an expansion in the coupling constant, which is the one we shall use in the following.² This is particularly convenient, since for the usual perturbation theory of QED we shall perform an expansion in the coupling constant in any case. Accordingly, we are able to avoid performing multiple expansions by taking terms of the expansion series of Seiberg-Witten map appropriately into account. We will also add a spinor field in the fundamental representation of the gauge field into the picture, thus inducing a mapping $(\Psi, A_\mu, \Lambda) \mapsto (\hat{\Psi}, \hat{A}_\mu, \hat{\Lambda})$.

The strategy in deriving the θ -exact Seiberg-Witten map, in a nutshell, is first to relate two gauge field theories in noncommutative spacetimes with infinitesimally differing noncommutativity parameter matrices, say θ and θ' , to each other in a gauge orbit preserving way, and then to integrate this relation from the origin $\theta_0 \equiv 0$ to some constant matrix θ_1 along a path in the space of 4×4 real-valued anti-symmetric matrices. Thus, let us have two noncommutative gauge field theories with spinor fields, denoted by $\mathcal{T}[\theta^{\mu\nu}, A_\mu, \Psi]$ and $\mathcal{T}'[\theta'^{\mu\nu}, A'_\mu, \Psi']$, where the arguments are the noncommutativity parameters, the gauge fields and the spinor fields, respectively. Let us also introduce the notation

$$\begin{aligned}\theta'^{\mu\nu} - \theta^{\mu\nu} &= \delta\theta^{\mu\nu} \quad , \\ A'_\mu - A_\mu &= a_\mu \quad , \\ \Psi' - \Psi &= \psi \quad .\end{aligned}\tag{1}$$

As prescribed, we assume that $\delta\theta^{\mu\nu}$ are infinitesimal, and that the fields depend smoothly on the noncommutativity parameters, so that a_μ , ψ and all their partial derivatives are also infinitesimal.

Let us now consider a mapping of the fields from $\mathcal{T}$ to $\mathcal{T}'$. We may think of the fields in $\mathcal{T}'$ as depending on the fields in $\mathcal{T}$ according to this mapping, namely³

$$A'_\mu \equiv A'_\mu(A) = A_\mu + a_\mu(A) \quad \text{and} \quad \Psi' \equiv \Psi'(\Psi, A) = \Psi + \psi(\Psi, A) \quad .\tag{2}$$

Now, we apply a gauge transformation in the theory $\mathcal{T}$ with a gauge transformation parameter Λ . For a noncommutative gauge field theory a gauge transformation is given

²Since an expansion in $\theta^{\mu\nu}$ may obscure the possible UV/IR mixing of the noncommutative theory, a θ -exact approach is essential.

³Precisely which arguments are needed here depends on, and is revealed by, the solutions found below, but for clarity they are already given here. Moreover, we have dropped the Lorentz indices of the arguments for simplicity, since it is clear how they are resumed.

by the formulae⁴

$$\begin{aligned}\delta_\Lambda A_\mu &= \partial_\mu \Lambda + ig[\Lambda \star A_\mu] \quad , \\ \delta_\Lambda \Psi &= ig\Lambda \star \Psi \quad ,\end{aligned}\tag{3}$$

where g is the coupling constant and the noncommutative $\star$ -product is the Moyal product defined as

$$f \star g = \exp \left[\frac{i}{2} \theta^{\mu\nu} \frac{\partial}{\partial x^\mu} \frac{\partial}{\partial y^\nu} \right] f(x)g(y) \Big|_{x=y} .\tag{4}$$

The fundamental requirement for the Seiberg-Witten map is that it should preserve the gauge equivalence classes of the theory, so that the transformation Λ in $\mathcal{T}$ corresponds to a gauge transformation

$$\Lambda' \equiv \Lambda'(\Lambda, A) = \Lambda + \lambda(\Lambda, A)\tag{5}$$

in $\mathcal{T}'$:

$$A'_\mu(A + \delta_\Lambda A) = A'_\mu(A) + \delta_{\Lambda'} A'_\mu(A) \quad ,\tag{6}$$

$$\Psi'(\Psi + \delta_\Lambda \Psi, A + \delta_\Lambda A) = \Psi'(\Psi, A) + \delta_{\Lambda'} \Psi'(\Psi, A) \quad .\tag{7}$$

By substituting the formulae (1) and (3) into (6) and (7), and using the relation

$$f \star' g = f e^{\frac{i}{2} \overleftarrow{\partial}_\mu (\theta + \delta\theta)^{\mu\nu} \overrightarrow{\partial}_\nu} g = f \star g + \frac{i}{2} \delta\theta^{\mu\nu} (\partial_\mu f) \star (\partial_\nu g) \quad ,\tag{8}$$

we arrive at the equations

$$\begin{aligned}& a_\mu(A + \delta_\Lambda A) - a_\mu(A) - \partial_\mu \lambda(\Lambda, A) - ig[\lambda(\Lambda, A) \star A_\mu] - ig[\Lambda \star a_\mu(A)] \\ &= -\frac{g}{2} \delta\theta^{\alpha\beta} \{ \partial_\alpha \Lambda \star \partial_\beta A_\mu \}\end{aligned}\tag{9}$$

and

$$\begin{aligned}& \psi(\Psi + \delta_\Lambda \Psi, A + \delta_\Lambda A) - \psi(\Psi, A) - ig\Lambda \star \psi(\Psi, A) - ig\lambda(\Lambda, A) \star \Psi \\ &= -\frac{g}{2} \delta\theta^{\alpha\beta} (\partial_\alpha \Lambda) \star (\partial_\beta \Psi)\end{aligned}\tag{10}$$

for λ , a_μ and ψ . As found by Seiberg and Witten in [2] (for $g \equiv 1$), the equation (9) is solved by

$$\begin{aligned}\lambda &= -\frac{g}{4} \delta\theta^{\alpha\beta} \{ A_\alpha \star \partial_\beta \Lambda \} \quad , \\ a_\mu &= -\frac{g}{4} \delta\theta^{\alpha\beta} \{ A_\alpha \star \partial_\beta A_\mu + F_{\beta\mu} \} \quad ,\end{aligned}\tag{11}$$

⁴We do not worry about gauge fixing here, since we shall ultimately fix it in the commutative QED to which we arrive in Section 3. Therefore, the Faddeev-Popov ghost fields are not needed.

where $F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu - ig[A_\mu \star A_\nu]$ is the field strength. Using (11), we find for the equation (10) the solution

$$\psi = -\frac{g}{2}\delta\theta^{\alpha\beta} \left[A_\alpha \star (\partial_\beta \Psi) + \frac{1}{2}(\partial_\beta A_\alpha) \star \Psi \right] . \quad (12)$$

As prescribed, the next step in constructing the θ -exact Seiberg-Witten map is to integrate these relations along a path in the space of real-valued anti-symmetric matrices to obtain a relation between gauge theories in a commutative spacetime and in a noncommutative one with finite noncommutativity parameters $\theta^{\mu\nu}$. There are certain ambiguities related to choosing a particular path, following from the observation that successive Seiberg-Witten maps do not commute in general, and thus there is an infinite number of free parameters related to the path fixing. Some but not all of these correspond to gauge transformations and field redefinitions, as explored in [5, 6]. However, for simplicity, we choose to consider a straight path⁵ $\gamma : [0, 1] \rightarrow \{\theta \in \mathbb{R}^{4 \times 4} | \theta \text{ antisymmetric}\}$ such that $\gamma(s) = s\theta_1$, where θ_1 is the constant matrix reached at $s = 1$. Let us denote the fields, now considered as dependent on the spacetime coordinates x^μ and the noncommutativity parameters $\theta^{\mu\nu}$, as $A_\mu(x; \theta)$ and $\Psi(x; \theta)$. Integrating the variation (9) along the straight path γ and using integration by parts, we get for the gauge field the equation

$$\begin{aligned} A_\mu(x; \theta_1) &= A_\mu(x; 0) + \lim_{y \rightarrow x} \left\{ -\frac{g\theta_1^{\alpha\beta}}{4} \frac{e^{\frac{i}{2}\theta^{\rho\sigma} \frac{\partial}{\partial x^\rho} \frac{\partial}{\partial y^\sigma}}}{\frac{i}{2}\theta_1^{\gamma\delta} \frac{\partial}{\partial x^\gamma} \frac{\partial}{\partial y^\delta}} \right. \\ &\quad \times \left[A_\alpha(x; \theta) (\partial_\beta A_\mu(y; \theta) + F_{\beta\mu}(y; \theta)) \right. \\ &\quad \left. \left. + (\partial_\beta A_\mu(x; \theta) + F_{\beta\mu}(x; \theta)) A_\alpha(y; \theta) \right] \right. \\ &\quad + \frac{g\theta_1^{\alpha\beta}}{4} \sum_{n=2}^{\infty} (-1)^n \frac{e^{\frac{i}{2}\theta^{\rho\sigma} \frac{\partial}{\partial x^\rho} \frac{\partial}{\partial y^\sigma}}}{\left(\frac{i}{2}\theta_1^{\gamma\delta} \frac{\partial}{\partial x^\gamma} \frac{\partial}{\partial y^\delta} \right)^n} \left(\prod_{k=2}^n \theta_1^{\alpha_k \beta_k} \frac{\delta}{\delta \theta^{\alpha_k \beta_k}} \right) \\ &\quad \times \left[A_\alpha(x; \theta) (\partial_\beta A_\mu(y; \theta) + F_{\beta\mu}(y; \theta)) \right. \\ &\quad \left. \left. + (\partial_\beta A_\mu(x; \theta) + F_{\beta\mu}(x; \theta)) A_\alpha(y; \theta) \right] \right\} \Bigg|_{\theta=0}^{\theta=\theta_1}, \quad (13) \end{aligned}$$

⁵This is the case considered also in Ref. [1].

and similarly for the spinor field the equation

$$\begin{aligned}
\Psi(x; \theta_1) = & \Psi(x; 0) + \lim_{y \rightarrow x} \left\{ - \frac{g\theta_1^{\alpha\beta}}{4} \frac{e^{\frac{i}{2}\theta^{\rho\sigma}}}{\frac{i}{2}\theta_1^{\gamma\delta}} \frac{\partial}{\partial x^\rho} \frac{\partial}{\partial y^\sigma} \right. \\
& \times \left[A_\alpha(x; \theta)(\partial_\beta \Psi(y; \theta)) + \frac{1}{2}(\partial_\beta A_\alpha(x; \theta))\Psi(y; \theta) \right] \\
& + \frac{g\theta_1^{\alpha\beta}}{4} \sum_{n=2}^{\infty} (-1)^n \frac{e^{\frac{i}{2}\theta^{\rho\sigma}}}{\left(\frac{i}{2}\theta_1^{\gamma\delta} \frac{\partial}{\partial x^\gamma} \frac{\partial}{\partial y^\delta}\right)^n} \left(\prod_{k=2}^n \theta_1^{\alpha_k \beta_k} \frac{\delta}{\delta \theta^{\alpha_k \beta_k}} \right) \\
& \times \left[A_\alpha(x; \theta)(\partial_\beta \Psi(y; \theta)) + \frac{1}{2}(\partial_\beta A_\alpha(x; \theta))\Psi(y; \theta) \right] \Bigg\}_{\theta=0}^{\theta=\theta_1},
\end{aligned} \tag{14}$$

which can be calculated iteratively in powers of the coupling constant g , since $\frac{\delta}{\delta \theta} A_\mu = \mathcal{O}(g)$ and $\frac{\delta}{\delta \theta} \Psi = \mathcal{O}(g)$, so the variations in the sums give terms of ever increasing powers in g .

3 NCQED via Seiberg-Witten map

We now turn to consider exclusively the gauge group $U(1)$, i.e., quantum electrodynamics (QED). We need to study the action of noncommutative QED,

$$\mathcal{S}_{\text{NCQED}} = \int d^4x \left[\hat{\bar{\Psi}} \star (i\not{\partial} - m) \hat{\Psi} - \frac{1}{4} \hat{F}_{\mu\nu} \star \hat{F}^{\mu\nu} - e \hat{\bar{\Psi}} \star \hat{A} \star \hat{\Psi} \right], \tag{15}$$

in terms of the ordinary fields up to the second order in the electromagnetic coupling constant e in order to catch all the second order contributions to the photon self-energy. These arise from the diagrams drawn in Fig. 1. Denoting the noncommutative fields by hats and dropping the lower index from θ_1 , we find the gauge field via the equation




Figure 1: Photon self-energy diagrams in the second order of e .

(13) up to second order in the coupling constant:

$$\begin{aligned}
\hat{A}_\mu^{(0)} &= A_\mu \quad , \\
\hat{A}_\mu^{(1)} &= -e \frac{\sin(\frac{1}{2}\partial_1 \wedge \partial_2)}{\frac{1}{2}\partial_1 \wedge \partial_2} \left[\eta_\mu^\beta \tilde{\partial}_2^\alpha - \frac{1}{2}\theta^{\alpha\beta} \partial_{2\mu} \right] A_\alpha(x_1) A_\beta(x_2) \Big|_{x_1=x_2} \\
&= -\frac{e}{2} \theta^{\alpha\beta} A_\alpha \star_1^s (2\partial_\beta A_\mu - \partial_\mu A_\beta) \quad , \\
\hat{A}_\mu^{(2)} &= \frac{e^2}{4} \left[\frac{\sin(\frac{1}{2}\partial_1 \wedge \partial_2)}{\frac{1}{2}\partial_1 \wedge \partial_2} \frac{\sin(\frac{1}{2}(\partial_1 + \partial_2) \wedge \partial_3)}{\frac{1}{2}(\partial_1 + \partial_2) \wedge \partial_3} \right. \\
&\quad \left. + \frac{\cos(\frac{1}{2}\partial_1 \wedge \partial_2) \cos(\frac{1}{2}(\partial_1 + \partial_2) \wedge \partial_3) - 1}{[\frac{1}{2}(\partial_1 + \partial_2) \wedge \partial_3]^2} \right] \\
&\quad \times \left\{ 2[2\tilde{\partial}_2^\alpha \tilde{\partial}_3^\beta - \theta^{\alpha\beta} \partial_2 \wedge \partial_3] \eta_\mu^\rho + 2[2(\tilde{\partial}_1 + \tilde{\partial}_2)^\rho \tilde{\partial}_2^\alpha - \theta^{\rho\alpha} \partial_1 \wedge \partial_2] \eta_\mu^\beta \right. \\
&\quad \left. + [(2\theta^{\beta\rho} \tilde{\partial}_2^\alpha - \theta^{\alpha\beta} (3\tilde{\partial}_1 + \tilde{\partial}_2)^\rho - 2\theta^{\rho\alpha} \tilde{\partial}_1^\beta) \partial_{2\mu} \right. \\
&\quad \left. - (\theta^{\alpha\beta} \tilde{\partial}_2^\rho + 2\theta^{\beta\rho} \tilde{\partial}_2^\alpha) \partial_{3\mu} \right\} A_\alpha(x_1) A_\beta(x_2) A_\rho(x_3) \Big|_{x_1=x_2=x_3} . \quad (16)
\end{aligned}$$

Similarly for the spinor field, via the equation (14), we find:

$$\begin{aligned}
\hat{\Psi}^{(0)} &= \Psi, \\
\hat{\Psi}^{(1)} &= -\frac{e}{2} \frac{e^{\frac{i}{2}\partial_1 \wedge \partial_2} - 1}{\frac{i}{2}\partial_1 \wedge \partial_2} \left(\frac{1}{2}\tilde{\partial}_1 + \tilde{\partial}_2 \right)^\alpha A_\alpha(x_1) \Psi(x_2) \Big|_{x_1=x_2} \\
&= -\frac{e}{2} \theta^{\alpha\beta} \left[A_\alpha \star_1 (\partial_\beta \Psi) + \frac{1}{2} (\partial_\beta A_\alpha) \star_1 \Psi \right], \\
\hat{\Psi}^{(2)} &= \frac{e^2}{4} \left\{ \left[\frac{\sin(\frac{1}{2}\partial_1 \wedge \partial_2)}{\frac{1}{2}\partial_1 \wedge \partial_2} \frac{e^{\frac{i}{2}(\partial_1 + \partial_2) \wedge \partial_3}}{\frac{i}{2}(\partial_1 + \partial_2) \wedge \partial_3} - \frac{\cos(\frac{1}{2}\partial_1 \wedge \partial_2) e^{\frac{i}{2}(\partial_1 + \partial_2) \wedge \partial_3} + 1}{[\frac{1}{2}(\partial_1 + \partial_2) \wedge \partial_3]^2} \right] \right. \\
&\quad \times \left[2\tilde{\partial}_2^\alpha \left(\frac{1}{2}\tilde{\partial}_1 + \frac{1}{2}\tilde{\partial}_2 + \tilde{\partial}_3 \right)^\beta + \theta^{\alpha\beta} \left(\frac{1}{2}\partial_1 + \partial_3 \right) \wedge \partial_2 \right] \\
&\quad + \left[\frac{e^{\frac{i}{2}\partial_1 \wedge \partial_3} - 1}{\frac{i}{2}\partial_1 \wedge \partial_3} \frac{e^{\frac{i}{2}\partial_2 \wedge (\partial_1 + \partial_3)}}{\frac{i}{2}\partial_2 \wedge (\partial_1 + \partial_3)} - \frac{e^{\frac{i}{2}\partial_1 \wedge \partial_3} e^{\frac{i}{2}\partial_2 \wedge (\partial_1 + \partial_3)} + 1}{[\frac{i}{2}\partial_2 \wedge (\partial_1 + \partial_3)]^2} \right] \\
&\quad \left. \times \left(\frac{1}{2}\tilde{\partial}_1 + \tilde{\partial}_3 \right)^\alpha \left(\tilde{\partial}_1 + \frac{1}{2}\tilde{\partial}_2 + \tilde{\partial}_3 \right)^\beta \right\} A_\alpha(x_1) A_\beta(x_2) \Psi(x_3) \Big|_{x_1=x_2=x_3}, \quad (17)
\end{aligned}$$

where we have introduced the notations $\tilde{u}^\mu := \theta^{\mu\nu} u_\nu$ and $u \wedge v := u_\mu \theta^{\mu\nu} v_\nu$ for any four-vectors u_μ, v_μ , and⁶

$$\begin{aligned}
f \star_1 g &:= \frac{e^{\frac{i}{2}\partial_1 \wedge \partial_2} - 1}{\frac{i}{2}\partial_1 \wedge \partial_2} f(x_1) g(x_2) \Big|_{x_1=x_2}, \\
f \star_1^s g &:= \frac{1}{2} \{f \star_1 g\} = \frac{\sin(\frac{1}{2}\partial_1 \wedge \partial_2)}{\frac{1}{2}\partial_1 \wedge \partial_2} f(x_1) g(x_2) \Big|_{x_1=x_2}. \quad (18)
\end{aligned}$$

Clearly, the terms tend to get more complicated at each order, which makes higher order calculations via θ -exact Seiberg-Witten map highly elaborate.

Since $(f \star g)^\dagger = g^\dagger \star f^\dagger$ for any functions (or, more generally, matrices) f and g , $\hat{\bar{\Psi}} \equiv \bar{\hat{\Psi}} = \hat{\Psi}^\dagger \gamma^0$. Substituting (16) and (17) into the action (15), we find the first order fermion-photon interaction term to be

$$\begin{aligned}
\mathcal{L}_{\bar{\Psi} A \Psi}^{(1)} &= -e \bar{\Psi} \star A \star \Psi \\
&\quad - \frac{e}{2} \theta^{\alpha\beta} \left[(\partial_\beta \bar{\Psi}) \star_1 A_\alpha + \frac{1}{2} \bar{\Psi} \star_1 (\partial_\beta A_\alpha) \right] \star (i\not{\partial} - m) \Psi \\
&\quad - \frac{e}{2} \theta^{\alpha\beta} \bar{\Psi} \star (i\not{\partial} - m) \left[A_\alpha \star_1 (\partial_\beta \Psi) + \frac{1}{2} (\partial_\beta A_\alpha) \star_1 \Psi \right]. \quad (19)
\end{aligned}$$

Similarly, from (16) we find the photon-photon interaction Lagrangian up to the first

⁶Notice that our notation for these so-called ‘generalized $\star$ -products’ differs from that used in [14, 15, 1]. This is an attempt to make the notation more systematic. The lower index denotes the times of integration of the $\star$ -product over the unit interval, and the upper index ‘s’ denotes symmetrization of the product with respect to its arguments.

order in e to be

$$\begin{aligned}\mathcal{L}_{A^3}^{(1)} = & -\frac{e}{4}\left\{\partial_\mu A_\nu - \partial_\nu A_\mu \star i[A^\mu \star A^\nu] \right. \\ & -\frac{1}{2}\theta^{\alpha\beta}\left[\partial^\mu\left(A_\alpha \star_1^s(2\partial_\beta A^\nu - \partial^\nu A_\beta)\right) \right. \\ & \left. \left. -\partial^\nu\left(A_\alpha \star_1^s(2\partial_\beta A^\mu - \partial^\mu A_\beta)\right)\right]\right\} .\end{aligned}\quad (20)$$

The second order contributions are considerably more complicated, but are obtained similarly by substituting the expressions (16) and (17) into the action (15) and picking up the terms with the factor e^2 . The Feynman diagram vertex functions arising from the first- and second-order parts of the action are given in Appendix A.

4 Photon self-energy corrections

Now, using the vertex function (32) from Appendix A, we find for the second order fermion loop correction to the photon propagator arising from the diagram (a) in Fig. 1 the form:

$$\begin{aligned}\Pi_{(a)}^{\alpha\beta}(k) = & -4e^2 \int \frac{d^4p}{(2\pi)^4} \\ & \times \left\{ T^{\alpha\beta} + \frac{i \sin(\frac{1}{4}p \wedge k)}{2 \frac{1}{4}p \wedge k} \left[\left(\tilde{p} - \frac{1}{2}\tilde{k}\right)^\alpha k_\rho T^{\rho\beta} e^{-\frac{i}{4}p \wedge k} \right. \right. \\ & \left. \left. - \left(\tilde{p} - \frac{1}{2}\tilde{k}\right)^\beta k_\rho T^{\rho\alpha} e^{\frac{i}{4}p \wedge k} \right] \right. \\ & \left. + \frac{1 \sin^2(\frac{1}{4}p \wedge k)}{4 \left(\frac{1}{4}p \wedge k\right)^2} \left(\tilde{p} - \frac{1}{2}\tilde{k}\right)^\alpha \left(\tilde{p} - \frac{1}{2}\tilde{k}\right)^\beta k_\rho k_\sigma T^{\rho\sigma} \right\} ,\end{aligned}\quad (21)$$

where

$$T^{\alpha\beta} := \frac{(p-k)^\alpha p^\beta + p^\alpha (p-k)^\beta + [m^2 - (p-k) \cdot p] \eta^{\alpha\beta}}{p^2(p-k)^2} ,\quad (22)$$

which is the only term we get in the commutative case. Here we see that we obtain the commutative result at the limit $\theta \rightarrow 0$ in contrast with the result of Schupp and You [1]. This follows from their use of the adjoint representation for the spinor field, leading one to antisymmetrize the $\star$ -products in the action, which gives rise to a sine phase factor for the vertex function. In our case of the fundamental representation for the fermions, on the other hand, one does not antisymmetrize the $\star$ -products, thus obtaining an exponential phase factor. The exponential factors cancel out upon the multiplication of complex conjugates arising from the two vertices of the diagram (a), leading to the usual commutative contribution, while the sine factors arising from the vertices in the adjoint representation do not cancel out upon multiplication. The first

term in (21) along with parts of the second and the third terms without phase factors can be interpreted as corresponding to the planar contribution of the diagram. The parts of the second and the third terms with nontrivial phase factors, on the other hand, correspond to the nonplanar part, which typically gives rise to UV/IR mixing.

To evaluate the parts of the second term in (21) with phase factors, we use the trick of Schupp and You [1] by expressing them as

$$\begin{aligned}
& 2 \int \frac{d^4 p}{(2\pi)^4} \frac{1}{\frac{1}{2}p \wedge k} \left[\left(\tilde{p} - \frac{1}{2}\tilde{k} \right)^\mu k_\rho T^{\rho\nu} e^{-\frac{i}{2}p \wedge k} + \left(\tilde{p} - \frac{1}{2}\tilde{k} \right)^\nu k_\rho T^{\rho\mu} e^{\frac{i}{2}p \wedge k} \right] \\
&= 2i \sum_{\lambda=\pm 1} \int d\lambda I^{\mu\nu}(k; \lambda) \quad , \quad (23)
\end{aligned}$$

where

$$I^{\mu\nu}(k; \lambda) = \int \frac{d^4 p}{(2\pi)^4} \left(\tilde{p} - \frac{1}{2}\tilde{k} \right)^\mu k_\rho T^{\rho\nu} e^{\frac{i}{2}\lambda p \wedge k} \quad . \quad (24)$$

By performing a Wick rotation $p^\mu = e_i^\mu \bar{p}^i$, where $e_i^\mu = \text{diag}(i, 1, 1, 1)$ and $\bar{p}^i$ is the Euclidean momentum, and using Schwinger parametrization

$$\frac{1}{\bar{p}^2 + m^2} = \int_0^\infty d\alpha e^{-\alpha(\bar{p}^2 + m^2)} \quad , \quad (25)$$

we get

$$\begin{aligned}
I^{\mu\nu}(k; \lambda) &= i e_i^\mu e_j^\nu \int \int_0^\infty d\alpha d\beta \int \frac{d^4 p}{(2\pi)^2} \left(\bar{p} - \frac{1}{2}\bar{k} \right)^i [(\bar{k}^2 - 2\bar{k} \cdot \bar{p})\bar{p}^j + (\bar{p}^2 + m^2)\bar{k}^j] \\
&\quad \times e^{-\alpha[(\bar{p}-\bar{k})^2 + m^2] - \beta[\bar{p}^2 + m^2] + \frac{i}{2}\lambda \bar{p} \cdot \bar{k}} \quad . \quad (26)
\end{aligned}$$

We may render the momentum integral Gaussian by applying the change of variables

$$\bar{q} := \bar{p} - \frac{\alpha}{\alpha + \beta} \bar{k} - \frac{i\lambda}{4(\alpha + \beta)} \bar{\bar{k}} \quad , \quad (27)$$

after which we can perform the integration over $\bar{q}$. Further, multiplying the integrand by

$$1 = \int_0^\infty dc \delta(c - \alpha - \beta) \quad , \quad (28)$$

changing the order of integrations, and applying the change of variables $\alpha = ca$, $\beta = cb$, we get

$$\begin{aligned}
I^{\mu\nu}(k; \lambda) &\approx \frac{i e_i^\mu e_j^\nu \bar{\theta}^{ik}}{(4\pi)^2} \int \int_0^1 da db \delta(1 - a - b) \int_0^\infty dc c^{-3} \\
&\quad \times \left[\left(\frac{i\lambda}{2} - \frac{i\lambda^3 \bar{k}^2}{64c} \right) \bar{\bar{k}}_k \bar{k}^j - \frac{i\lambda}{4} \bar{\bar{k}}_k \bar{k}^j \right] \\
&\quad \times e^{-c(ab\bar{k}^2 + m^2) - \frac{\lambda^2}{16c} \bar{\bar{k}}^2} \quad , \quad (29)
\end{aligned}$$

where the less IR-divergent terms are dropped out.⁷ The dependence on a and b drops out, and the integrals over them give unity. The integral over λ is now straightforward to perform. Moreover, the integral over c can be performed and expressed for small k using the properties of modified Bessel functions $K_r(x, y)$ [17]:

$$\int_0^\infty dc \, c^{-r-1} e^{-xc-y/c} = 2 \left(\frac{x}{y} \right)^{\frac{r}{2}} K_r [2\sqrt{xy}] \quad , \text{ where } \operatorname{Re}[x], \operatorname{Re}[y] > 0 \quad ,$$

$$\text{and } K_r(z) \approx \frac{\Gamma(r)}{2} \left(\frac{2}{z} \right)^r \quad , \text{ when } 0 < z \ll \sqrt{r+1} \quad . \quad (30)$$

We get for small $\tilde{k}^2 \ll m^{-2}$ accordingly

$$i\Pi_{(a)\text{np}}^{\mu\nu}(k) \approx \frac{8e^2}{\pi^2} \frac{\tilde{k}^\mu \tilde{k}^\nu}{\tilde{k}^4} + \frac{4e^2}{\pi^2} \frac{\tilde{k}^\mu k^\nu + k^\mu \tilde{k}^\nu}{\tilde{k}^4} \quad . \quad (31)$$

The first term here is similar to the IR-divergent terms found in the usual formulation of NCQED and by Schupp and You [1]. The second term gives another quadratic IR-divergence, which is gauge variant, and therefore should be cancelled, when all the second order contributions in the coupling constant are taken into account.

Having found the gauge invariant IR-divergence in (31), we proceed to confirm the absence of canceling terms. The calculations, though more elaborate, follow precisely the same scheme as the one above. To make them manageable we only consider contributions of the form $a\tilde{k}^\mu \tilde{k}^\nu$, where a is a scalar quantity. The third term in (21) and the other second order contributions coming from the diagrams (b), (c) and (d) also give rise to quadratic divergencies of the form $c\tilde{k}^\mu \tilde{k}^\nu / \tilde{k}^4$, where c is a constant. For all of the contributions we find $c > 0$, and thus they cannot cancel the IR-divergence of (31). Hence we conclude that the UV/IR mixing problem persists in the noncommutative QED formulated here via θ -exact Seiberg-Witten map.

5 Conclusions and remarks

We have found that UV/IR mixing is present in the photon self-energy corrections of noncommutative QED defined via θ -exact Seiberg-Witten map for a straight path in θ -space. The result further demonstrates that UV/IR mixing is a generic property of noncommutative quantum field theories, and is not cured in general by the approach via Seiberg-Witten map, contrary to some claims previously made in the literature.

A question remains open, though, whether the result holds generally for all possible integration paths in θ -space. It is not ruled out that by modifying the integration path one could get rid of the divergence, although on mathematical grounds this seems

⁷Here we have to take into account the following integration over c , where $c \sim \tilde{k}^2$. The integration over λ does not affect the relative powers of divergence.

unlikely, at least, for paths obtainable from the straight one by smooth deformations. Of course, answering the question properly requires a rigorous analysis, which we postpone to a future study.

In the case of a scalar field theory in noncommutative spacetime, as for example in [16], the destruction of UV/IR mixing by θ -expansion becomes immediately obvious. In [16] there is also given a satisfactory explanation for the mixing as a direct result of the infinite nonlocality of $\star$ -product, and thus it is deeply rooted in the very definition of noncommutativity of spacetime. It therefore seems unlikely that the resulting divergencies could be made vanish, at least, without modifying the theory itself by introducing new terms in the Lagrangian, which suppress the contributions of the IR sector. This has been done for a noncommutative scalar field theory in [18, 19, 20] and for noncommutative QED in [21, 22]. For some other attempts, see [23, 24, 25]. Reducing the nonlocality of the noncommutative field theories to a finite range is also an option which has been preliminarily exploited in [26, 27].

All in all, what ever the direction, more work is to be done before we are to overcome the obstacles arising from the nonlocality in noncommutative quantum field theories.

Acknowledgements

We would like to thank Masud Chaichian for many helpful discussions. The support of the Academy of Finland under the Projects No. 121720 and 127626 is gratefully acknowledged.

A Vertex functions

The momenta are everywhere incoming, p_i 's for fermions and k_i 's for photons.

For the first order vertex functions we get the expressions:

$$V_{\Psi A \Psi}^{\mu}(p_1, p_2) = -ie\gamma^{\mu}e^{\frac{i}{2}p_1 \wedge p_2} - \frac{ie}{2}(\tilde{p}_1 - \tilde{p}_2)^{\mu}(\not{p}_1 + \not{p}_2)\frac{e^{\frac{i}{2}p_1 \wedge p_2} - 1}{p_1 \wedge p_2} \quad , \quad (32)$$

$$\begin{aligned} & V_{A^3}^{\mu_1 \mu_2 \mu_3}(k_1, k_2, k_3) \\ = & 2e \sin\left(\frac{k_1 \wedge k_2}{2}\right) \\ & \times \left\{ (k_1 - k_2)^{\mu_3} \eta^{\mu_1 \mu_2} + \frac{1}{\frac{1}{2}k_1 \wedge k_2} \left[(k_1^{\mu_1} k_1^{\alpha} - k_1^2 \eta^{\mu_1 \alpha})(2\tilde{k}_3^{\mu_2} \eta_{\alpha}^{\mu_3} - k_{3\alpha} \theta^{\mu_2 \mu_3}) \right] \right\} \\ & + \{symm.\} \quad , \quad (33) \end{aligned}$$

where $\{symm.\}$ denotes terms symmetrizing the previous contributions with respect to the photons (k_i, μ_i) .

For the second order vertex functions we similarly find

$$\begin{aligned}
& V_{\tilde{\Psi}A^2\Psi}^{\mu_1\mu_2}(k_1, k_2, p_1, p_2) \\
= & -\frac{ie^2}{4} \left\{ \left[\left(\frac{\sin(\frac{1}{2}k_1 \wedge k_2)}{\frac{1}{2}k_1 \wedge k_2} \frac{e^{\frac{i}{2}p_1 \wedge p_2}}{\frac{i}{2}p_1 \wedge p_2} - \frac{\cos(\frac{1}{2}k_1 \wedge k_2)e^{\frac{i}{2}p_1 \wedge p_2} - 1}{(\frac{i}{2}p_1 \wedge p_2)^2} \right) \right. \right. \\
& \times \left((\tilde{p}_1 - \tilde{p}_2)^{\mu_1} \tilde{k}_1^{\mu_2} - \frac{1}{2} \theta^{\mu_1\mu_2} (p_1 - p_2) \wedge k_1 \right) \\
& + \left(\frac{e^{\frac{i}{2}p_2 \wedge k_1}}{\frac{i}{2}p_2 \wedge k_1} \frac{e^{\frac{i}{2}k_2 \wedge p_1} - 1}{\frac{i}{2}k_2 \wedge p_1} - \frac{e^{\frac{i}{2}p_2 \wedge k_1} e^{\frac{i}{2}k_2 \wedge p_1} - 1}{(\frac{i}{2}p_2 \wedge k_1)^2} \right) \\
& \times \frac{1}{4} (\tilde{p}_1 - \tilde{p}_2 + \tilde{k}_2)^{\mu_1} (\tilde{p}_1 - \tilde{p}_2 - \tilde{k}_1)^{\mu_2} \left. \right] (\not{p}_2 - m) \\
& - \left[\left(\frac{\sin(\frac{1}{2}k_1 \wedge k_2)}{\frac{1}{2}k_1 \wedge k_2} \frac{e^{\frac{i}{2}p_1 \wedge p_2}}{\frac{i}{2}p_1 \wedge p_2} - \frac{\cos(\frac{1}{2}k_1 \wedge k_2)e^{\frac{i}{2}p_1 \wedge p_2} - 1}{(\frac{i}{2}p_1 \wedge p_2)^2} \right) \right. \\
& \times \left((\tilde{p}_2 - \tilde{p}_1)^{\mu_2} \tilde{k}_2^{\mu_1} - \frac{1}{2} \theta^{\mu_2\mu_1} (p_2 - p_1) \wedge k_2 \right) \\
& + \left(\frac{e^{\frac{i}{2}p_2 \wedge k_1}}{\frac{i}{2}p_2 \wedge k_1} \frac{e^{\frac{i}{2}k_2 \wedge p_1} - 1}{\frac{i}{2}k_2 \wedge p_1} - \frac{e^{\frac{i}{2}p_2 \wedge k_1} e^{\frac{i}{2}k_2 \wedge p_1} - 1}{(\frac{i}{2}p_2 \wedge k_1)^2} \right) \\
& \times \frac{1}{4} (\tilde{p}_2 - \tilde{p}_1 + \tilde{k}_1)^{\mu_2} (\tilde{p}_2 - \tilde{p}_1 - \tilde{k}_2)^{\mu_1} \left. \right] (\not{p}_1 + m) \\
& + \frac{1}{4} \frac{e^{\frac{i}{2}k_1 \wedge p_1} - 1}{\frac{i}{2}k_1 \wedge p_1} \frac{e^{\frac{i}{2}k_2 \wedge p_2} - 1}{\frac{i}{2}k_2 \wedge p_2} (\tilde{p}_1 - \tilde{p}_2 - \tilde{k}_2)^{\mu_1} (\tilde{p}_2 - \tilde{p}_1 - \tilde{k}_1)^{\mu_2} (\not{p}_1 + \not{p}_2 + m) \\
& + i \frac{e^{\frac{i}{2}k_1 \wedge p_1} - 1}{\frac{i}{2}k_1 \wedge p_1} e^{\frac{i}{2}p_2 \wedge k_2} (\tilde{p}_1 - \tilde{p}_2 - \tilde{k}_2)^{\mu_1} \gamma^{\mu_2} \\
& + i \frac{e^{\frac{i}{2}k_2 \wedge p_2} - 1}{\frac{i}{2}k_2 \wedge p_2} e^{\frac{i}{2}p_1 \wedge k_1} (\tilde{p}_2 - \tilde{p}_1 - \tilde{k}_1)^{\mu_2} \gamma^{\mu_1} \\
& + 4i \frac{\sin(\frac{1}{2}k_1 \wedge k_2)}{\frac{1}{2}k_1 \wedge k_2} e^{\frac{i}{2}p_1 \wedge p_2} (\tilde{k}_2^{\mu_1} \gamma^{\mu_2} - \frac{1}{2} \theta^{\mu_1\mu_2} \not{p}_2) \left. \right\} \\
& + \{symm.\} \quad , \tag{34}
\end{aligned}$$

$$\begin{aligned}
& V_{A^4}^{\mu_1\mu_2\mu_3\mu_4}(k_1, k_2, k_3, k_4) \\
= & -e^2 \left\{ \left[\frac{\sin(\frac{1}{2}k_1 \wedge k_2)}{\frac{1}{2}k_1 \wedge k_2} \frac{\sin(\frac{1}{2}k_3 \wedge k_4)}{\frac{1}{2}k_3 \wedge k_4} - \frac{\cos(\frac{1}{2}k_1 \wedge k_2) \cos(\frac{1}{2}k_3 \wedge k_4) - 1}{(\frac{1}{2}k_3 \wedge k_4)^2} \right] \right. \\
& \times \left[\left(\tilde{k}_2^{\mu_1} \tilde{k}_3^{\mu_2} - \frac{1}{2} \theta^{\mu_1\mu_2} k_2 \wedge k_3 \right) \eta_\alpha^{\mu_3} \right. \\
& + \left((\tilde{k}_1 + \tilde{k}_2)^{\mu_3} \tilde{k}_2^{\mu_1} - \frac{1}{2} \theta^{\mu_3\mu_1} k_1 \wedge k_2 \right) \eta_\alpha^{\mu_2} \\
& + \frac{1}{2} \left(\theta^{\mu_2\mu_3} \tilde{k}_2^{\mu_1} + \theta^{\mu_1\mu_3} \tilde{k}_1^{\mu_2} - \frac{1}{2} \theta^{\mu_1\mu_2} (3\tilde{k}_1 + \tilde{k}_2)^{\mu_3} \right) k_{2\alpha} \\
& \left. - \frac{1}{2} \left(\theta^{\mu_2\mu_3} \tilde{k}_2^{\mu_1} + \frac{1}{2} \theta^{\mu_1\mu_2} \tilde{k}_2^{\mu_3} \right) k_{3\alpha} \right] \\
& \times \left(k_4^2 \eta^{\alpha\mu_4} - k_4^\alpha k_4^{\mu_4} \right) \\
& - \frac{1}{2} \frac{\sin(\frac{1}{2}k_1 \wedge k_2)}{\frac{1}{2}k_1 \wedge k_2} \frac{\sin(\frac{1}{2}k_3 \wedge k_4)}{\frac{1}{2}k_3 \wedge k_4} \left(\eta_\alpha^{\mu_2} \tilde{k}_2^{\mu_1} - \frac{1}{2} \theta^{\mu_1\mu_2} k_{2\alpha} \right) \\
& \times \left(\eta_\beta^{\mu_4} \tilde{k}_4^{\mu_3} - \frac{1}{2} \theta^{\mu_3\mu_4} k_{4\beta} \right) \left((k_1 + k_2)^2 \eta^{\alpha\beta} - (k_1 + k_2)^\alpha (k_1 + k_2)^\beta \right) \\
& + \frac{\sin(\frac{1}{2}k_1 \wedge k_2)}{\frac{1}{2}k_1 \wedge k_2} \sin(\frac{1}{2}k_3 \wedge k_4) \left(\eta_\alpha^{\mu_2} \tilde{k}_2^{\mu_1} - \frac{1}{2} \theta^{\mu_1\mu_2} k_{2\alpha} \right) \\
& \times \left(\eta^{\mu_3\mu_4} k_4^\alpha - \eta^{\alpha\mu_4} k_4^{\mu_3} \right) \\
& + \sin(\frac{1}{2}k_1 \wedge k_2) \frac{\sin(\frac{1}{2}k_3 \wedge k_4)}{\frac{1}{2}k_3 \wedge k_4} \left(\eta_\alpha^{\mu_4} \tilde{k}_4^{\mu_3} - \frac{1}{2} \theta^{\mu_3\mu_4} k_{4\alpha} \right) \\
& \times \left(\eta_\alpha^{\mu_2} (k_3 + k_4)^{\mu_1} - \eta_\alpha^{\mu_1} (k_3 + k_4)^{\mu_2} \right) \\
& \left. + \sin(\frac{1}{2}k_1 \wedge k_2) \sin(\frac{1}{2}k_3 \wedge k_4) \eta^{\mu_1\mu_2} \eta^{\mu_3\mu_4} \right\} \\
& + \{symm.\} \quad , \tag{35}
\end{aligned}$$

where again $\{symm.\}$ denotes terms symmetrizing the previous contributions with respect to the photons (k_i, μ_i) .

References

- [1] Peter Schupp and Jiangyang You, *UV/IR mixing in noncommutative QED defined by Seiberg-Witten map*, JHEP **0808** 107 (2008), arXiv:0807.4886 [hep-th].
- [2] Nathan Seiberg and Edward Witten, *String Theory and Noncommutative Geometry*, JHEP **9909** 032 (1999), arXiv:hep-th/9908142.

- [3] Sergio Doplicher, Klaus Fredenhagen and John E. Roberts, *The Quantum Structure of Spacetime at the Planck Scale and Quantum Fields*, Commun. Math. Phys. **172** 187 (1995), arXiv:hep-th/0303037.
- [4] Daniel N. Blaschke, Erwin Kronberger, Arnold Rofner, Manfred Schweda, René I.P. Sedmik and Michael Wohlgenannt, *On the Problem of Renormalizability in Non-Commutative Gauge Field Models - A Critical Review*, arXiv:0908.0467 [hep-th].
- [5] Tsuguhiko Asakawa and Isao Kiskimoto, *Comments on Gauge Equivalence in Noncommutative Geometry*, JHEP **9911** 024 (1999), arXiv:hep-th/9909139.
- [6] Bing Suo, Pei Wang and Liu Zhao, *Ambiguities of the Seiberg-Witten map in the presence of matter fields*, Commun. Theor. Phys. **37** 571 (2002), arXiv:hep-th/0111006.
- [7] M. Chaichian, P. Prešnajder, M.M. Sheikh-Jabbari and A. Tureanu, *Noncommutative Gauge Field Theories: A No-Go Theorem*, Phys. Lett. **B526** 132 (2002), arXiv:hep-th/0107037.
- [8] M. Chaichian, P. Prešnajder, M.M. Sheikh-Jabbari and A. Tureanu, *Can Seiberg-Witten Map Bypass Noncommutative Gauge Theory No-Go Theorem?*, Phys. Lett. **B 683** 55 (2010), arXiv:0907.2646 [hep-th].
- [9] Andreas Bichl, Jesper Grimstrup, Harald Grosse, Lukas Popp, Manfred Schweda and Raimar Wulkenhaar, *Renormalization of the noncommutative photon self-energy to all orders via Seiberg-Witten map*, JHEP **0106** 013 (2001), arXiv:hep-th/0104097.
- [10] M.M. Sheikh-Jabbari, *Open Strings in a B-field Background as Electric Dipoles*, Phys. Lett. **B 455** 129 (1999), arXiv:hep-th/9901080.
- [11] Daniela Bigatti and Leonard Susskind, *Magnetic fields, branes and noncommutative geometry*, Phys. Rev. **D 62** 066004 (2000), arXiv:hep-th/9908056.
- [12] Zheng Yin: *A Note On Space Noncommutativity*, Phys. Lett. **B 466** 234 (1999), arXiv:hep-th/9908152.
- [13] L. Álvarez-Gaumé and J.L.F. Barbon, *Non-linear Vacuum Phenomena in Non-commutative QED*, Int. J. Mod. Phys. **A 16** 1123 (2001), arXiv:hep-th/0006209.
- [14] Mohammad R. Garousi, *Non-commutative world-volume interactions on D-brane and Dirac-Born-Infeld action*, Nucl. Phys. **B 579** 209 (2000), arXiv:hep-th/9909214.
- [15] Thomas Mehen and Mark B. Wise, *Generalized $\star$ -Products, Wilson Lines and the Solution of the Seiberg-Witten Equations*, JHEP **0012** 008 (2000), arXiv:hep-th/0010204.

- [16] Shiraz Minwalla, Mark Van Raamsdonk and Nathan Seiberg, *Noncommutative Perturbative Dynamics*, JHEP **0002** 020 (2000), arXiv:hep-th/9912072.
- [17] W. Magnus, F. Oberhettinger and R.P. Soni, *Formulas and Theorems for the Special Functions of Mathematical Physics*, 3rd Ed., Springer-Verlag, Berlin Heidelberg 1966.
- [18] Harald Grosse and Raimar Wulkenhaar, *Renormalization of ϕ^4 -theory on non-commutative $\mathbb{R}^4$ in the matrix base*, Commun. Math. Phys. **256** 305 (2005), arXiv:hep-th/0401128.
- [19] Harald Grosse and Michael Wohlgenannt, *Noncommutative QFT and Renormalization*, J. Phys. Conf. Ser. **53** (2006) 764, arXiv:hep-th/0607208.
- [20] R. Gurau, J. Magnen, V. Rivasseau and A. Tanasa, *A translation-invariant renormalizable non-commutative scalar model*, Commun. Math. Phys. **287** 275 (2009), arXiv:0802.0791.
- [21] Daniel N. Blaschke, Arnold Rofner, Manfred Schweda and Rene I.P. Sedmik, *One-Loop Calculations for a Translation Invariant Non-Commutative Gauge Model*, Eur. Phys. J. **C 62** 433 (2009), arXiv:0901.1681 [hep-th].
- [22] L.C.Q. Vilar, O.S. Ventura, D.G. Tedesco, V.E.R. Lemes, *Renormalizable Non-commutative $U(1)$ Gauge Theory Without IR/UV Mixing*, arXiv:0902.2956 [hep-th].
- [23] P. Fischer and V. Putz, *No UV/IR Mixing in Unitary Space-Time Noncommutative Field Theory*, Eur. Phys. J. **C 32** 269 (2004), arXiv:hep-th/0306099.
- [24] S. Denk and M. Schweda, *Time ordered perturbation theory for non-local interactions: Applications to NCQFT*, JHEP **09** 032 (2003), arXiv:hep-th/0306101.
- [25] Stefan Denk, Volkmar Putz and Michael Wohlgenannt, *Consistent Construction of Perturbation Theory on Noncommutative Spaces*, Eur. Phys. J. **C 45** (2006) 263, arXiv:hep-th/0402229.
- [26] D. Bahns and S. Waldmann, *Locally Noncommutative Space-Times*, Rev. Math. Phys. **19** 273 (2007), arXiv:math/0607745.
- [27] M. Långvik and A. Zahabi, *On Finite Noncommutativity in Quantum Field Theory*, arXiv:1002.0956 [hep-th].